

Probing internal excitation of trapped $O^+(^4S, ^2D, ^2P)$ ions by reaction with N_2



R. Plašil¹, A. Kovalenko¹, T. D. Tran¹, S. Rednyk¹, Š. Roučka¹, P. Dohnal¹, J. Glosík¹

¹ Department of Surface and Plasma Science, Faculty of Mathematics and Physics, Charles University in Prague, Prague, Czech Republic

When the atomic oxygen cation O^+ takes part as a reactant or a product in experimental studies, electronically excited states may play a significant role. For investigation of ion molecular reactions, a cryo-cooled radiofrequency ion trap is used. Molecular nitrogen was chosen as a monitor gas for probing a presence of long-lived excited $O^+(^2D, ^2P)$ in an ensemble of ions confined in the radiofrequency trap. The fraction of electronically excited ions was determined. The rate coefficient of the reaction of $O^+(^4S)$, the electronic ground state, with nitrogen molecule was measured at low temperatures. The reaction rate coefficient at 61 K was evaluated as $(7.5 \pm 1.5) \times 10^{-12} \text{ cm}^3 \text{ s}^{-1}$. It confirms an increase of the coefficient at low temperatures.

Introduction

The atomic oxygen cation O^+ plays a significant role in numerous environments such as planetary ionospheres or technological plasmas. The existence of long-lived $O^+(^2D, ^2P)$ metastable states may be important in many situations. For states $^2D_{3/2}$ and $^2D_{5/2}$ the lifetimes are in the order of hours and for states $^2P_{1/2}$ and $^2P_{3/2}$ in seconds [1]. The excited states carry electronic energy of 3.3 eV and 5.0 eV, respectively. The only exoergic channel for reaction of O^+ in ground state with N_2 is formation of NO^+ (1). The charge transfer (2) is endoergic.



For the excited states $O^+(^2D, ^2P)$ both reactions are exoergic and for reaction (2) the rate coefficient is close to Langevin rate in the order of $10^{-10} \text{ cm}^3 \text{ s}^{-1}$ [1, 2]. The charge transfer (2) from excited O^+ ion to N_2 is dominant channel.

We used these reactions as a probe for excited states of O^+ ions confined in the radiofrequency trap. In addition we confirmed steep increase of the rate coefficient of the reaction (2) with decreasing temperature [1, 3].

Experiment

A 22-pole radiofrequency ion trap was used for this study. It was placed on a cryo-cooler in an ultra-high vacuum system. The measuring procedure was based on iterative filling of the trap with a well-defined number of primary ions O^+ . They react with N_2 and the contents of the trap were analysed after chosen times by means of a quadrupole mass spectrometer with micro-channel plate detector. Additional details may be found in reference [4] and references therein. The O^+ ions were produced in separated ion source by electron impact from N_2O

molecule to reduce the ratio of excited states in comparison with a production from O_2 .

From the number of N_2^+ we may evaluate a fraction of excited states of O^+ as $(5 \pm 1)\%$ for electron impact energy $E_e = 75 \text{ eV}$ and $(10 \pm 2)\%$ for 145 eV, see Figure 1.




Fig. 1: Evolution of normalized number of ions in rf trap at temperature 61 K. From the increase of NO^+ we may determine the rate coefficient of the reaction (1) $(7.5 \pm 1.5) \times 10^{-12} \text{ cm}^3 \text{ s}^{-1}$. The number densities of the N_2 reactant and helium buffer gas are indicated in the figure.

We thank the TU Chemnitz, the DFG, prof. D. Gerlich for lending us the apparatus and Czech Science Foundation grant 17-18067S.

References

- [1] J. Glosík *et al.*, J. Phys. B: Atom. Molec. Phys. **11** (1978) 3365.
- [2] B.R. Rowe *et al.*, J. Chem. Phys. **73** (1980) 194.
- [3] J-L. Le Garrec *et al.*, Chem. Phys. Lett. **372** (2003) 485.
- [4] D. Gerlich and G. Borodi, Faraday Discussions **142** (2009) 57.