

Electron/molecular-cation collisions in cold plasmas: super-excited states at "zero" energy

J. Zs. Mezei^{1,2,3,4}, F. Colboc¹, Y. Moulane^{1,5,6}, N. Pop^{1,7}, S. Niyonzima^{1,8}, M. D. Epée Epée^{1,9},
O. Motapon^{1,9,10}, D. A. Little¹¹, F. Iacob¹², R. Boata¹³, V. Laporta^{1,11,14}, R. Celiberto^{14,15},
K. Chakrabarti^{1,16}, E. Jehin⁶, Z. Benkhaldoun⁵, K. Hassouni², D. Benredjem³, A. Faure¹⁷,
A. Bultel¹⁸, J. Tennyson¹¹, I. F. Schneider^{1,3}

¹Laboratoire Ondes et Milieux Complexes, CNRS, Université du Havre, Le Havre, France

²Laboratoire des Sciences des Procédés et des Matériaux, CNRS, Université Paris 13, Villetaneuse, France

³Laboratoire Aimé Cotton, CNRS, ENS Cachan and Université Paris-Sud, Orsay, France

⁴Institute of Nuclear Research of the Hungarian Academy of Sciences, Debrecen, Hungary

⁵Oukaimeden Observatory, High Energy Physics & Astrophysics Lab., Cadi Ayyad Univ., Marrakech, Morocco

⁶Institut d'Astrophysique et de Géophysique, Liège, Belgium

⁷Department of Physical Foundations of Engineering, Politehnica University Timișoara, Timișoara, Romania

⁸Département de Physique, Université du Burundi, Bujumbura, Burundi

⁹Department of Physics, Faculty of Sciences, University of Douala, Douala, Cameroon

¹⁰University of Maroua, Faculty of Science, Maroua, Cameroon

¹¹Department of Physics and Astronomy, University College London, United Kingdom

¹²Physics Faculty, West University of Timișoara, Timișoara, Romania

¹³Astronomical Institute of the Romanian Academy, Timisoara Astronomical Observatory, Timișoara, Romania

¹⁴Istituto di Nanotecnologia, CNR, Bari, Italy

¹⁵Dipartimento di Ingegneria Civile, Ambientale, del Territorio, Edile e di Chimica, Politecnico di Bari, Italy

¹⁶Department of Mathematics, Scottish Church College, Calcutta, India

¹⁷Université de Grenoble Alpes, CNRS, Inst. de Planétologie et d'Astrophysique de Grenoble, Grenoble, France


¹⁸Laboratoire CORIA, CNRS, Université de Rouen, Saint Etienne de Rouvray, France

We will discuss the role of excited states in cold ionized media, as resonances in reactive collisions, and as initial states of the target in non-equilibrium regime.

Dissociative recombination (DR) [1,2]:



the major recombination process in low pressure ionized gases, is often notably fast at "zero" (i.e. low) energy of the incident electrons, since it occurs via super-excited molecular states singly- (AB^*) or doubly- (AB^{**}) excited, embedded in the ionization continuum of the target ion. This process is competed by ro-vibrational and dissociative excitation:



The use of the Multichannel Quantum Defect Theory [3] resulted in accurate state-to-state cross sections and rate coefficients, displaying a resonant character and a strong dependence on the target state. These features are particularly important for the collisional-radiative modeling of the cold ionized gases – Fig. 1 – in various environments - interstellar space, comets, planetary ionospheres, shock-waves in the entries of spacecrafts, ionic propulsion devices, industrial and edge fusion plasmas - and containing various cations - H_2^+ , BeH^+ , BF^+ [2,4], N_2^+ [5], CO^+ [6], SH^+ , ArH^+ , H_3^+ [7], BF_2^+ , etc.




Fig.1. Maxwell rate coefficients for the Dissociative Recombination (DR) of H_2^+ : dependence on the vibrational temperature of the ions.

References

- [1] I. F. Schneider, O. Dulieu, J. Robert, *Proc. of DR2013: The 9th Int. Conf. on Dissociative Recombination: Theory, Experiment and Applications*, Paris, July 7-12, 2013, EPJ Web of Conferences **84** (2015).
- [2] N. Pop *et al*, poster contribution to this conference.
- [3] O. Motapon *et al*, *Phys. Rev. A* **90** (2014) 012706.
- [4] V. Laporta *et al*, *Pl. Phys. Contr. Fusion* **59** (2017) 045008.
- [5] D. A. Little *et al*, *Phys. Rev. A* **90** (2014) 052705.
- [6] J. Zs. Mezei *et al.*, *Pl. Sour. Sci. Tech.* **24** (2015) 035005.
- [7] I. F. Schneider *et al*, *Phys. Rev. A* **86** (2012) 062706.