

Transport properties of hot dense plasmas

S.K. Kodanova¹, T.S. Ramazanov¹, M.K. Issanova¹, E.E. Shokparbayeva¹

¹*IETP, Al-Farabi Kazakh National University, Almaty, Kazakhstan*

In this work the transport properties of non-isothermal dense plasmas were studied. Transport processes in dense plasmas were studied on the basis of the effective potentials using the Coulomb logarithm. These potentials take into consideration long-range multi-particle screening effects and short-range quantum-mechanical effects in two-temperature plasmas. The obtained results were compared with the theoretical works of other authors and with the results of MD simulations.

Investigation of transport properties of the dense plasma is a great importance for plasma physics, as well as for the problems of inertial confinement fusion (ICF), warm dense matter driven by heavy ion beams [1]. Experimental investigation of dense nonideal plasmas based on using of a shock wave compression, a high-power laser and an ion accelerator devices [1-2].

In presented work transport properties of hot dense, non-isothermal plasma are considered. One of the important values describing the transport coefficients of deuterium-tritium plasma is the Coulomb logarithm. The Coulomb logarithm is obtained on the basis of effective potentials. These interaction potentials take into consideration long-range many particle screening effects as well as short-range quantum-mechanical effects [3].

The Coulomb logarithm is determined by the center of mass scattering angle θ_c [4-5]:

$$\lambda_{\alpha\beta} = \frac{1}{b_{\perp}^2} \int_0^{b_{\max}} \sin^2\left(\frac{\theta_c}{2}\right) b db, \quad (1)$$

$$\theta_c = \pi - 2b \int_{r_0}^{\infty} \frac{dr}{r^2} \left(1 - \frac{\Phi_{\alpha\beta}(r)}{E_c} - \frac{b^2}{r^2} \right)^{1/2}, \quad (2)$$

where $E_c = \frac{1}{2} m_{\alpha\beta} v^2$ is the energy of the center of mass, $m_{\alpha\beta} = m_{\alpha} m_{\beta} / (m_{\alpha} + m_{\beta})$ is the reduced mass of the particles of kinds α and β (electron and ion); $b_{\perp} = Z_{\alpha} Z_{\beta} e^2 / (m_{\alpha\beta} v^2)$, $b_{\min} = \max\{b_{\perp}, \lambda_{\alpha\beta}\}$ describes the minimum impact parameter, where is $\lambda_{\alpha\beta} = \hbar / \sqrt{2\pi n_{\alpha\beta} k_B T}$ is the thermal de Broglie wavelength.

For inertial confinement fusion applications, we have calculated diffusion and viscosity of deuterium-tritium plasma for density $\rho = 5 \text{ g/cm}^3$ and temperatures ranging from 2 to 10 eV using the Coulomb Logarithm based on effective potentials. Fig.1 show a comparison of the calculated data on diffusion and viscosity in a DT plasma with the

theoretical results of other authors [6] such as finite-temperature Kohn-Sham density-functional theory molecular dynamics (QMD) and orbital-free molecular dynamics (OFMD).




Fig. 1. The diffusion (a) and viscosity (b) coefficients for the DT plasma as function of temperature.

References

- [1] V.E. Fortov et al. Physics-Uspekhi **51** (2) (2008).
- [2] N.A. Tahir et al. Phys. Plasmas **18** (2011) 032704.
- [3] T.S. Ramazanov, Zh.A. Moldabekov, M.T. Gabdullin. Phys.Rev. E **92** (2015) 023104.
- [4] M.K. Issanova, S.K. Kodanova, T.S. Ramazanov, et al. LPB **34** (2016) 457-466.
- [5] T.S. Ramazanov, et al. Contrib. Plasma Phys. **56** (5) (2016) 425-431.
- [6] J.D. Kress, J.S. Cohen, D.A. Horner, et al. Phys. Rev. E **82** (2010) 036404.